

ServSafe® Food Handler Guide for Food Banking



Disclaimer

©2014 National Restaurant Association Educational Foundation. All rights reserved.

The information presented in this publication is provided for informational purposes only and is not intended to provide legal advice or establish standards of reasonable behavior. Operators who develop food safety-related policies and procedures are urged to obtain the advice and guidance of legal counsel. Although National Restaurant Association Solutions, LLC (NRA Solutions) endeavors to include accurate and current information compiled from sources believed to be reliable, NRA Solutions, and its licensor, the National Restaurant Association Educational Foundation (NRAEF), affiliates, distributors, and agents (hereafter collectively referred to as "NRAS") make no representations or warranties as to the accuracy, currency, or completeness of the information. No responsibility is assumed or implied by the NRAS for any damage or loss resulting from inaccuracies or omissions or any actions taken or not taken based on the content of this publication.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Sections 107 and 108 of the 1976 United States Copyright Act, without prior written permission of the publisher.

Request to use or reproduce material from this publication should be directed to:

Copyright Permissions National Restaurant Association Solutions 175 West Jackson Boulevard, Suite 1500 Chicago, IL 60604-2814 Email: Permissions@restaurant.org

ServSafe® and the ServSafe logo are registered trademarks of the National Restaurant Association Educational Foundation, and used under license by National Restaurant Association Solutions, LLC, a wholly owned subsidiary of the National Restaurant Association. National Restaurant Association and the arc design are trademarks of the National Restaurant Association.

Food Handler Guide for Food Banking 10 Pack–EG6RFA10

ISBN 978-1-58280-320-3

Printed in the USA 10 9 8 7 6 5 4 3 2 1

ACKNOWLEDGEMENTS

The development of the ServSafe Food Handler Guide for Food Banking would not have been possible without the expertise of our many advisors and manuscript reviewers. Thank you to the following people and organizations for their time, effort, and dedication to creating this guide.

Nicole Acinelli, Food Lifeline Maurice Casey, Vermont Food Bank Megan DeMers, San Antonio Food Bank Kris DeSmedt, Northern Illinois Food Bank Karen Erickson, Northeast Iowa Food Bank Barbara Gehlen, Second Harvest Food Bank of Santa Clara & San Mateo Counties Amy Gerhart-Wackerhagen, Food Bank for the Heartland Jeff Gutierrez, Montana Food Bank Network Gordon Hahn and Team, Northern Illinois Food Bank Micheline Hynes, Tarrant Area Food Bank Stuart Iseminger, M.S.W., Lakeview Pantry Linda Lundquist, Second Harvest Food Bank of Southern Wisconsin Jesse Sharrard, Greater Pittsburgh Food Bank Jane MacDonald, Cary Gilkey, and Team, Loaves and Fishes Ralph Maltese, Second Harvest Food Bank of Santa Clara & San Mateo Counties Sam Michaud, Good Shepherd Food Bank Mark Mitchell, Akron/Canton Food Bank Kimberly Molnar, Second Harvest Food Bank of Middle Tennessee Steve Randolph, Oregon Food Bank Amythst Shipman, Food Lifeline Jessica Simpson, Eastern Illinois Food Bank Anthony Stover, Food Lifeline Jim Thomas. Food Bank of the Albemarle Dani Turk, Food Lifeline Julia Williamson, Oregon Food Bank Gina Wilson, Second Harvest Food Bank of Southern Wisconsin Jordan Vernoy, Feeding America National Office

TABLE OF CONTENTS

Food Safety Is Important

How Food Becomes Unsafe	1-1
Your Role in Keeping Food Safe	1-3
Understanding Food Allergies	1-4
Preventing Food Allergen Contamination	1-5

Good Personal Hygiene

How and When to Wash Your Hands	2-1
Where to Wash Your Hands	2-5
Other Hand-Care Guidelines	2-6
What to Wear	2-8
Other Important Practices	2-9

Receiving and Storing Food Safely

Controlling Time and Temperature During Receiving	3-1
Inspecting Food During Receiving to Make Sure It Is Safe	3-3
Storing Food Safely	3-5

Evaluating, Repacking, and Transporting Food Safely

Evaluating the Condition of Food	4-1
Repacking Food	4-6
Loading and Distributing Food Safely	4-8

Cleaning and Sanitizing

How and When to Clean and Sanitize	5-1
How to Manually Clean and Sanitize Tools and Utensils	5-6
Handling Garbage	5-8
Spotting Pests	5-9

Apply Your Knowledge Answers

A-1

T-1

Food Safety Is Important



- How food becomes unsafe
- Your role in keeping food safe
- Understanding food allergies
- Preventing food allergen contamination



Unsafe Temperatures

An independent trucking company was forced to discard several thousand pounds of food products being transported in one of its delivery trucks. A random check by officials found that meat and other perishables on the truck were being held at unsafe temperatures. The food was discarded because officials feared it could cause a foodborne illness.

How FOOD BECOMES UNSAFE

People who receive food from a food bank or agency trust you to help keep them safe. One big concern is foodborne illness. A foodborne illness is a disease that is transmitted to people through food.

Many hazards can make food unsafe and cause a foodborne illness. Some come from the air, water, soil, and animals. Some come from dirty surfaces. Others happen when people don't handle food correctly. Foodborne illness is almost always preventable. You just need to know how to handle food safely.

Hazards from the Environment

Three types of hazards make food unsafe. They are: biological hazards, physical hazards, and chemical hazards.



Biological hazards

These are tiny forms of life that you can't see, taste, or smell. They include:

- Bacteria
- Viruses
- Parasites
- Fungi

These forms of life are everywhere. Sometimes they are harmless. But some cause illness. They are called pathogens.



Physical hazards

Some physical hazards occur naturally in food, such as bones in fillets and fruit pits.

Usually physical hazards occur when objects fall into food. Some examples include:

- Jewelry
- Glass
- Brittle plastics
- Ceramics
- Bandages



Chemical hazards

Chemicals in your food bank or agency can contaminate food. These include:

- Cleaners
- Sanitizers
- Pesticides

How People Make Food Unsafe

Sometimes the things people do can make food unsafe. Once you understand what the dangers are, it is easy to avoid them. These four practices can make food unsafe.



Poor personal hygiene

Transferring pathogens from your body to food.

This is the number one cause of foodborne-illness outbreaks.

Do Nor Use Jai gi

Crosscontamination

Transferring pathogens from one surface or food to another.



Time-temperature abuse

Letting food stay too long at temperatures that are good for pathogen growth.



Poor cleaning and sanitizing

Transferring pathogens from incorrectly cleaned surfaces to food.

Apply Your Knowledge

What Do You Think?

Write an \mathbb{X} next to the problem for each action listed below.

 Leaving raw chicken breasts on a pallet in a loading area that is not refrigerated A Time-temperature abuse B Poor personal hygiene C Cross-contamination D Poor cleaning and sanitizing 	 Sneezing on food A Time-temperature abuse B Poor personal hygiene C Cross-contamination D Poor cleaning and sanitizing
3 Keeping produce that raw meat juices have leaked on	Scraping off food from an otherwise clean food-storage container
A Time-temperature abuse	A Time-temperature abuse
B Poor personal hygiene	B Poor personal hygiene
C Cross-contamination	C Cross-contamination
D Poor cleaning and sanitizing	D Poor cleaning and sanitizing

YOUR ROLE IN KEEPING FOOD SAFE

You play an important role in keeping food safe. Knowing the hazards to food safety is just the start. The next step is to make sure you work safely.

It is easy but critical that you follow these practices.



Practice good personal hygiene

- DON'T transfer pathogens from your body to food.
- You'll learn more about this in Chapter 2.



Control the time and temperature of food

- DON'T let food stay too long at temperatures that are good for pathogen growth.
- You'll learn more about this in Chapter 3.



Prevent cross-contamination

- DON'T transfer pathogens from one food to another.
- DON'T transfer pathogens from one surface to another.
- You'll learn more about this in Chapters 3 and 4.



Clean and sanitize surfaces correctly

- Keep everything clean.
- Clean and sanitize anything that touches food.
- You'll learn more about this in Chapter 5.

UNDERSTANDING FOOD ALLERGIES

Some people are allergic to certain types of food. The tiniest speck of a food they are allergic to can make them sick or even cause death. You must keep this food away from these people.

When a food item containing an allergen comes in contact with another food item and their proteins mix, it is called cross-contact. This can be dangerous for people with food allergies. Food should be stored and handled in a way that prevents cross-contact.

The Most Common Food Allergens

The proteins that cause allergic reactions are called allergens. Many different food items can cause allergic reactions. But just eight food items cause most reactions. These are called the Big Eight.

You need to be aware of the Big Eight and the products that contain them. Here are the Big Eight allergens.





Soy

Milk



Eggs



• Fish, such as bass, flounder, and cod



 Tree nuts, such as almonds, walnuts, and pecans



Peanuts



 Crustacean shellfish, such as crab, lobster, and shrimp



• Wheat

PREVENTING FOOD ALLERGEN CONTAMINATION

Labeling is the most important way to help keep people with food allergies safe. You will learn more about correct labeling in Chapter 3. The following practices can also prevent food-allergen contamination.

Prevent Cross-Contact

You must make sure that food containing allergens does not come in contact with other food.



- Clean and sanitize all surfaces that have come in contact with food containing an allergen.
- Inspect food packaging to ensure no leaks or spills have occurred that can cause cross-contact.
- When possible, store food with allergens separately from allergenfree products. **DO NOT** store food containing allergens above allergen-free food. Use dedicated pallets and bins for products containing allergens.
- Wash your hands and change gloves after handling a food containing allergens and before handling an allergen-free food.



The Most Common Food Allergens

Write an X next to each food that is a common food allergen or contains one.

1	Теа	7	Potatoes	13	Squash and eggplant
2	Cod	8	Mushrooms	14	Soybeans
3	Wheat flour	9	Tomatoes	15	Rice and rice
4	Melons	10	Pecan pie		products
5	Peanut butter	11	Citrus fruit	16	Omelet
6	Crab legs	12	Green peppers	17	Vanilla ice cream

Cleaning Up Food Spills

Spills can easily lead to cross-contact. Follow these steps to prevent this.



- Immediately isolate any spilled food containing an allergen from other food products.
- Inspect surrounding products to make sure they have not come in contact with the spilled food.
- Dispose of any open products that have come in contact with the spilled food. If the food is in packaging that can be safely cleaned and sanitized, it may not need to be discarded.
- Clean and sanitize the area.

Good Personal Hygiene

111



- How and when to wash your hands
- Where to wash your hands
- Other hand-care guidelines
- What to wear
- Other important practices



Hepatitis A Scare

Hepatitis A vaccinations were offered to thousands of clients who visited a food pantry on the Gulf Coast of the United States. The vaccinations were offered by the local regulatory authority after a staff member at the operation tested positive for hepatitis A, exposing clients to the virus.

The identified staff member was responsible for packaging produce. He was excluded from work until approved to return by a physician and the regulatory authority.

The local regulatory authority also worked with the pantry's director and supervisors to ensure they had all of the correct processes in place to protect clients and staff. Ensuring that staff members reported illness to managers was at the top of the list.

How and When to Wash Your Hands

Handwashing is the most important part of personal hygiene. It is important because your hands can transfer pathogens to food. So it is critical that you wash your hands correctly and at the correct times.

How to Wash Your Hands

Correct handwashing only takes about 20 seconds. Follow these steps.



Wet hands and arms.

Use running water as hot as you can comfortably stand.



Apply soap. Apply enough to build up a good lather.



Scrub hands and arms vigorously. Scrub them for 10 to 15 seconds. Clean under fingernails and between fingers.



Rinse hands and arms thoroughly. Use running warm water.



Dry hands and arms.

DO NOT use your apron or any part of your clothing. Use a single-use paper towel or hand dryer.

After Washing Your Hands

Once your hands are clean, you don't want to contaminate them again. Follow these guidelines.



• Use a paper towei to turn off the faucet.



• Use a paper towel to open the restroom door.

When to Wash Your Hands

It is easy to contaminate your hands while doing everyday activities. And contaminated hands can spread pathogens.

To keep food safe, wash your hands before you start work and after any of these activities.



Using the restroom.

Failure to do this has caused many foodborne illnesses



• Handling raw meat, poultry, or seafood (before and after).





- Touching your hair, face, or body.
- Touching clothing or aprons.



• Taking out garbage.



 Sneezing, coughing, or using a tissue.



• Handling chemicals that can make food unsafe.



• Smoking.



Chewing gum or tobacco.



 Eating or drinking.



Before putting on gloves at the start of a new task.



 Touching anything else that may contaminate your hands.
 Examples include dirty equipment, work surfaces, and cloths.



 Leaving and returning to the food-handling area.

Using Hand Antiseptics

Some food banks or agencies may provide hand antiscptics, commonly called hand sanitizers, to reduce pathogens on hands. Follow these steps if you use hand antiseptics.



- **NEVER** use a hand antiseptic instead of washing your hands.
- Use an antiseptic after you wash your hands.
- Wait for the antiseptic to dry before touching food or equipment, and before putting on gloves.
- Follow the manufacturer's directions for using a hand antiseptic.

Apply Your Knowledge Check Your Handwashing Knowledge

1 Ryan washed his hands before handling food. Write an X next to each error that Ryan made.

A	He applied hand sanitizer before washing his hands	E	He rinsed his hands under running warm water
В	He wet his hands using very hot water	F	He dried his hands with his clean apron
C	He applied enough soap to develop a good lather	G	He turned off the faucet with a paper towel
D	He scrubbed his hands with soap for five seconds	Н	He opened the door with his clean hand s
2 Write	an $ imes$ next to each situation where yo	ou must wa	sh your hands.
A	After handling packages of raw chic	ken	
В	B Before putting on gloves at the start of a task		
C	Before handling chemicals		
D	After taking out garbage		
3 Write	an $ imes$ next to the situation where you	ı should use	e a hand antiseptic.
A	When you can't wash your hands		
В	Before washing your hands		
C After washing your hands			
D	When you can't dry your hands		

WHERE TO WASH YOUR HANDS

Your food bank or agency should have specific sinks for handwashing. You must use these sinks correctly to make handwashing effective.

Use a Handwashing Sink

Wash your hands only in a designated handwashing sink.





- **DO NOT** use handwashing sinks for other things.
- **NEVER** dump dirty water in them.
- **NEVER** prep food in them.
- **NEVER** wash tools or equipment in them.



Make sure handwashing sinks are easy to get to and are not blocked.

• **NEVER** stack food, equipment, or supplies in them or in front of them.

Stocking the Handwashing Sink



A stocked sink should have:

- Hot and cold running water
- Liquid soap
- Single-use paper towels or a hand dryer
- Garbage container

If these items aren't stocked, tell your director or supervisor.

OTHER HAND-CARE GUIDELINES

It takes more than just clean hands to maintain good personal hygiene. You also must use gloves correctly and care for your hands and nails.

Use Gloves Correctly

Gloves can help prevent the spread of pathogens. Here is the correct way to use them.

Use the correct gloves



• Only use singleuse gloves when handling food.



- ×
- NEVER rinse, wash, or reuse gloves.



- Make sure the gloves fit your hands. They should not be too tight or too loose.
- Never blow into a glove, or roll them to make them easier to put on.
- Wash your hands before putting on gloves when starting a new task.

Change gloves when necessary



 As soon as they become dirty or torn.





 After handling raw meat, seafood, or poultry and before handling ready-to-eat food.



- Before beginning a different task.
- Before or after handling any food with a known food allergen
- After an interruption, such as taking a phone call.

Hands and Nails

Follow these guidelines to maintain your hands and nails.





• Keep fingernails short and clean. Long fingernails can be hard to keep clean.

- If you will be handling exposed food, you must use single-use gloves if you are wearing:
 - Nail polish
 - ° False fingernails
- Nail polish can hide dirt under nails and may flake off into exposed food.



Infected wounds can contain bacteria. To keep food safe, make sure wounds are correctly covered.

- **Hands or wrist:** Cover with a bandage or finger cot that prevents fluid from leaking out. Then place a single-use glove over the cover.
- Arms: Completely cover with a bandage that prevents fluids from leaking out.
- Body: Cover with a dry, durable, tight-fitting bandage.



Keeping It Safe

- Alicia is a volunteer repackaging bulk cereal. Write an X beside everything that Alicia has done wrong.
 - A Washed her hands before putting on gloves
 - B Rinsed her gloves when they became too dirty from working with the cereal
 - C Began repackaging rice after repackaging cereal without changing gloves
 - __ D Washed her hands and changed gloves when she noticed a small tear in a glove she was wearing

- 2 Write an \times next to each unsafe practice.
 - A Washing hands and putting on new gloves after handling an allergen
 - B Putting the same gloves back on after finishing one task and starting a new task
- _____ C Prepping food wearing fake nails without gloves
 - ____ D Working with an infected, unbandaged cut

WHAT TO WEAR

If you fail to bathe or you wear dirty clothes while working with food, you can make people sick. Dirty hair, skin, and clothing may carry pathogens that can cause foodborne illness. Hygiene matters, so bathe daily. And follow the guidelines below.



Hair covering

Always wear a clean hat or other hair covering when:

- Repackaging food
- Working in a clean room
- Working in areas used to clean utensils and equipment

Men with beards and mustaches should also wear a beard restraint.

Clothing and aprons

- Wear clean clothes whenever working with or around food.
- Remove and throw away disposable aprons when leaving clean rooms or food-handling areas.
- Store personal belongings away from food.

Jewelry

Jewelry can contain bacteria and other pathogens. It can also fall off into food. Remove jewelry from hands and arms before handling exposed food or when working in or around food-handling areas. **DO NOT** wear:

- Rings, except for a plain band
- Bracelets, including medical bracelets
- Watches

Your director or supervisor may also ask you to remove other jewelry.

Apply Your Knowledge

It's What You Wear

1) Write an 🔀 next to each unsafe practice.

- _____1 Wearing a dirty shirt
- 2 Wearing nail polish without wearing gloves
- _____3 Wearing a baseball cap while packaging food
- _____4 Wearing a watch

- 5 Removing a disposable apron and putting it back on after returning to the clean room
- 6 Wearing a bandage on your finger under your gloves
- ____7 Cleaning and sanitizing food scoops without wearing a hat or other hair covering
- _____8 Wearing the same pants several days in a row until they get dirty

OTHER IMPORTANT PRACTICES

Following correct hygiene practices helps keep you and everyone else safe. Follow the quidelines below.

Eating, Drinking, Smoking, and Chewing Gum or Tobacco

Saliva contains pathogens that can cause a foodborne illness. Eating, drinking, smoking, and chewing gum or tobacco can contaminate equipment and your hands with saliva. Cigarette butts and other items can also get into food.



- NEVER eat, drink, smoke, or chew gum or tobacco in food-handling and storage areas.
- Only do these things in designated areas.

What to Do if You Are Sick

If you are sick, you could spread pathogens to food and equipment. This could make your clients and your coworkers sick. Don't let this happen.



Tell your director or supervisor if you have been diagnosed with a foodborne illness. You must also tell your director or supervisor if you have any of these symptoms:

- Vomiting
- Diarrhea
- Jaundice (yellowing of skin and eyes)
- Sore throat with a fever

pply Your Knowledg

eral scents let a let a failer e let a

Reporting Illnesses

 \bigcirc Write an imes next to the symptoms you must report to your director or supervisor.

_____1 Vomiting

_____ 4 Being very tired

- _____ 3 Sore throat with a fever
- _____5 Diarrhea
- _____6 Headache

For answers, please turn to the Answer Key.

Receiving and Storing Food Safely



- Controlling time and temperature during receiving
- Inspecting food during receiving to make sure it is safe
- Storing food safely



Salmonella Outbreak Threatens Food Banks

Food banks across the U.S. were forced to destroy thousands of pounds of food products containing peanuts. The culprit—a deadly strain of *Salmonella*, which killed nine people and sickened more than 700 people nationwide. No known cases of illness were attributed to the food distributed by food banks, thus pointing to the importance of following recall notices.

CONTROLLING TIME AND TEMPERATURE DURING RECEIVING

The temperature of food must be controlled from the minute it's picked up from the donor or food bank to the minute it's handed to the client.



- That includes the time food spends in the warehouse, on the truck, and at the agency.
- Think of this as a "cold chain" that, if broken, can result in foodborne illness.
- The cold chain is broken whenever food is held at temperatures between 41°F and 135°F (5°C and 57°C). This is the temperature danger zone.
- Fortunately, there are several ways to control temperature and prevent a "break" in the cold chain.

Types of Thermometers

A thermometer is the most important tool for controlling the temperature of food. Three types of thermometers are commonly used in food banks and agencies.



Infrared thermometers

- Only measure surface temperatures of food and equipment
- Display temperatures instantly
- Prevent cross-contamination and damage to food because they do not touch it



Thermocouples and other digital thermometers

- Do not need to be inserted deep into food to get an accurate reading
- Display temperatures quickly
- Many come with other probes for checking liquids, air temperature, and equipment



Bimetallic stemmed thermometers

- Must be inserted in food from the tip of the thermometer stem to the dimple to get an accurate reading
- Require at least 15 seconds before a final reading can be taken
- Can be easily calibrated at the food bank or agency

The Correct Way to Calibrate a Thermometer

Thermometers must be calibrated (adjusted) regularly to make sure the readings are correct.

- Digital thermometers may need to be sent back to the manufacturer to be calibrated. If not, follow the directions from the manufacturer to calibrate them.
- Infrared thermometers will need to be sent back to the manufacturer to be calibrated.

Follow these steps to calibrate a bimetallic stemmed thermometer.



Fill a large container with crushed ice and tap water.

• Stir the mixture well.



Put the thermometer stem or probe into the ice water.

- Make sure the sensing area is under water.
- DO NOT let the probe touch the container.
- Wait 30 seconds or until the indicator stops.



Adjust the thermometer so it reads 32°F (0°C).

- Hold the calibration nut with a wrench or other tool.
- Rotate the thermometer head until it reads 32°F (0°C).

poly Your nowledge Ice-Point Calibration Method

Write an imes next to the correct answer.

- 1 How long should you wait to read a bimetallic stemmed thermometer after placing it in the ice water?
- _____A 5 seconds
- _____B 7 seconds
- _____C 10 seconds
 - ____ D 30 seconds

What temperature should a thermometer be adjusted to after placing it in the ice water?

A	0°F (-18°C)
В	10°F (-12°C)
C	22°F (-6°C)

____ D 32°F (0°C)

INSPECTING FOOD DURING RECEIVING TO MAKE SURE IT IS SAFE

Checking food when you receive it will help make sure it is safe. This starts with inspecting delivery trucks for the correct temperature and signs of contamination. It continues with checking the temperature and quality of food and looking for signs of contamination.

Inspecting the Delivery Vehicle

It is very important that you inspect the delivery vehicle before unloading it. Any problems should be documented. Here's what you should look for.



• Overall condition of the vehicle



• Signs of pests in the product or vehicle



 Condition of the product



Presence of objectionable product (e.g., homemade product)



 Door locks and seals that are in place and functioning



- Correct truck temperature
- Check the temperature recording device or use an infrared thermometer



Reject the food delivery if the delivery vehicle or delivery containers:

- Do not meet acceptable standards
- Contain pests
- Are excessively dirty

Checking the Temperature of Food

Perishable food must be received at the correct temperature in order for it to be safe. Reject any food that does not meet these temperatures.

Type of food	Temperature
Refrigerated food	41°F (5°C) or lower.
Frozen food	Frozen solid. Recommend 0°F (-18°C) or lower.
Cut produce	41°F (5°C) or lower.
Whole produce	Dry storage temperatures. For more information see Feeding America's produce handling guidelines.

Apply Your Knowledge

Is the Delivery Safe?

1 Write an imes next to each unsafe situation.

- _____A The door lock on a food delivery truck is broken.
- _____ B Mice are spotted inside a food delivery truck.
 - _____ C A delivery truck is transporting food at 38°F (3°C).
 - D A case of home-canned jelly is found inside a delivery truck.

Apply Your Is It the Correct Temperature?

1 Write an X next to each item that was not received at the correct temperature.

- _____ A Frozen meat received at 38°F (3°C)
- _____ B Bags of cut lettuce received at 50°F (10°C)
- _____ C Bags of cut melons received at 45°F (7°C)
- _____D Milk received at 50°F (10°C)

STORING FOOD SAFELY

Food can easily be contaminated if it is not stored correctly. It can also become unsafe if it is allowed to stay in the temperature danger zone. The good news is that there are things you can do to prevent this.

General Storage Guidelines

Here are some important steps you must take to keep food safe when storing it.



Store refrigerated food at 41°F (5°C) or lower.

This includes cut produce.





• Keep frozen food frozen solid.

> The recommended temperature is 0°F (-18°C) or lower.



Store food only in designated foodstorage areas.

- Store food at least six inches (15 cm) off the floor.
- Store food away from walls.



 Store ready-to-eat food above raw meat, seafood, and poultry.

Do not store these items on the same shelf or pallet.



- Store food only in containers made for food.
- Wrap or cover food before storing it.

Is It Stored Correctly?

f 1 Draw an X through each food item that has not been stored correctly.



Labeling Food for Storage

All food given to clients must contain a label that includes specific information about the product. See Feeding America labeling guidelines for more information. While not required by law, food should also be labeled with the receiving date. This will ensure that products can be correctly rotated.



pply Your

nowledge

Put the receiving date on food and products.

- Dates should be placed in these locations:
 - ^o On the individual container or case
 - On pallet tags
- Make sure the dates are listed on a permanent part of the packaging.

Understanding Food Product Dating

It can be difficult to understand the dates listed on a food label. Here is an explanation of common labeling terms.



Packing or manufacturing date

- Used by manufacturers for tracking and recalls.
- Not an expiration date.



Sell-by date

- This is a quality date.
- It tells the store how long to display the product for sale.
- The product is still safe to eat past this date.





Best-by or best if used by date

- This is a quality date.
- It tells clients the date by which the product should be eaten for best flavor or quality.
- The product is still safe to eat past this date.

Use-by or expiration date

- This is the last date recommended for the product while at peak quality.
- The product is still safe to eat past this date.

Determining the Shelf-Life for Refrigerated Food in Storage

Most storage dates on food are not related to food safety. So it can be difficult to decide how long food can be safely stored. Here are some general guidelines to follow.



- If the food has a sell-by date or no date at all, you should freeze or distribute the food following Feeding America guidelines.
- Freezing food will keep it safe for long periods of time. But quality will suffer if frozen food is stored too long. Follow Feeding America guidelines for recommended storage times.

Handling Recalled Food

Sometimes manufacturers or government agencies will issue recalls for food. This may happen when food has been contaminated. It can also happen when food allergens have not been identified on the label.



- Talk to your director or supervisor for how to handle recalls in your food bank or agency.
- Feeding America issues recall notices for all FDA and USDA issued recalls.

Rotating Food in Storage

You must make sure that the oldest food in storage is used first. This will reduce food waste and keep food safe. Food rotation should be based on either the expiration date or the delivery date of the food.

Follow the first-expired, first-out (FEFO) method of rotation if the food has a use-by or expiration date.



1 Check the use-by or expiration date.

2 Store food that will expire first in front of items that will expire later.

3 Use the food stored in front first.

Follow the first-in, first-out (FIFO) method of rotation if the food does not have a use-by or expiration date.



Check the delivery date.



Store food with the earliest delivery date in front of food with later dates.



Use the food stored in front first.

Has It Been Rotated Correctly?

1 The date is 2014. Circle the storage shelf that has the cans stored correctly.



pply Your

Knowledge



Notes



Evaluating, Repacking, and Transporting Food Safely



- Evaluating the condition of food
- Repackaging food
- Loading and distributing food safely



Botulism Scare

Workers at a large food bank on the west coast were recently faced with sorting through thousands of pounds of donated canned goods. They were trying to keep food containing deadly bacteria from reaching the hungry. The effort came after five people in three states came down with botulism after eating some of the product. A national recall was issued for over 90 products including 27 brands of canned chili, stew, hash, and gravy. Fortunately the food bank was able to pull more than 300 cans before they were distributed to those in need.

EVALUATING THE CONDITION OF FOOD

Not all of the products you receive will be ready for distribution. Many products will require a closer look to see if they are usable. Each product will be evaluated differently. Never assume that the product has already been inspected and is safe for use. It is everyone's responsibility to inspect food for safety.

Checking Cans for Defects

Canned food that is damaged or defective can cause a foodborne illness. Here are some common can defects. Discard cans if they have these problems.



Checking Jars and Bottles for Defects

Food in jars and bottles must be checked just as carefully as canned food. Here are some common defects. Discard jars and bottles of food if they have these problems. Never accept homemade food.



Checking Commercially Packaged Dry Food for Defects

Commercially packaged dry food must be checked just as carefully as other food.



- Dry food containers include bags, boxes, sacks, or plastic-wrapped items.
- Some dry food is packaged in a single package. Other food is double-packaged with the food contained in an inner package.
- Food packaged this way includes flour, cereals, pasta, bread, and rice.

Discard commercially packaged dry food if it has an unusual appearance, is not in its original package, or has these problems.



- Unlabeled or not correctly labeled
- Label is unreadable
- No code dates
- nplete ncake Mix
- Signs of pests
 - ° Gnaw marks
 - ^o Droppings
 - Insects (dead or alive)
 - Pin-sized holes in packaging



 Double-packaged food with inner package that is damaged, wet, and/or stained



 Single-packaged food that is exposed due to packaging that is open, ripped or torn, punctured or cut, and/or split at seams



Keep It or Toss It?

1 Draw an imes through each food item that should be rejected.



Checking Produce for Defects

There are many ways that produce will show signs of spoilage. You should be able to recognize the obvious signs as well as those that indicate the produce will spoil quickly. Discard produce if it has these problems.



- Mold.
- Decay or rot.
- Bad odors.





 Signs of insects (live insects, insect bodies, or insect eggs).



- Severe bruises.
- Skin not intact.

Bruises provide a way for pathogens to get inside produce.

- Cut produce not at 41°F (5°C) or lower.
- No code dates.



• Produce repacked in packaging not safe for use with food.



 \bigcirc Draw an X through each food item that should be rejected.



Checking Over-the-Counter (OTC) Medications for Defects

Check with your director or supervisor to see if OTC medications can be received in your food bank or agency. Never accept products containing ephedrine or pseudoephedrine. Here's what to look for when evaluating OTC medications.



Product packaging

- Reject medications with these problems:
- Broken tamper-resistant seals.
- Damaged packaging.



Expiration and use-by dates.

Reject medications with these problems:

• Past the use-by date.

24.4	Nurfax
110 miles	-
•	The same particular
1. C	- Maren
	and the second second
9	
-	1. 100101
	tion in the local division of the local divi
2	C STREET
£.	-
	1

Product labels

Reject medications with these problems:

- Label is not intact or readable.
- Label does not include necessary information, including common name of drug, list of ingredients, guantity of contents, instructions for correct use, recommended dose, and storage requirements, if anv.

Apply Your Knowledge

Is It Acceptable?

1 Draw an \times through each item that should be rejected.







REPACKING FOOD

Many times food is received in bulk and must be repacked in quantities that are the right size for clients. There are strict conditions for repacking food. In general, it must be performed in a designated clean room. A clean room is a separate enclosed room dedicated to bulk repacking.



- It must be designed to the highest standards of sanitation.
- It must prevent as many environmental contaminants as possible.
- Interior surfaces must be durable and easy to clean.
- It must be built to commercial foodservice construction standards.
- It must meet local health regulations.

TCS food should only be repacked in a clean room that can maintain a temperature of 55°F (13°C) or lower.

Guidelines for Repacking Food

Follow these guidelines when repacking exposed food, especially food that is ready-to-eat.



 Practice correct personal hygiene before handling food.

(See Chapter 2)



Clean and sanitize the room before repacking food.



- Clean and sanitize equipment used to repack food.
 - Before and after use
 - When changing to a new product or between allergens
- with equipment made for food.



Only repack one type of product at a time.



Store clean and sanitized equipment so it doesn't become contaminated.



Only handle food

Labeling Repacked Food

All repackaged food must contain a label that includes specific information about the product.



What Did They Do Wrong?

1 Write an x next to each situation where a mistake was made.



pply Your

nowledge





LOADING AND DISTRIBUTING FOOD SAFELY

Food can become unsafe if it's not loaded or distributed correctly. Start by making sure the delivery vehicle is ready for handling food. You also need to be careful when getting food ready for a delivery and when loading the delivery vehicle.

Preparing Delivery Vehicles before Transporting Food

Delivery vehicles must be inspected and maintained to protect food from contamination. That's true whether they are private vehicles or commercial trucks. Here's what you should do to keep food safe.



- Clean the inside of vehicles at least once per week or as often as necessary.
- Make sure vehicles are free of pests, such as insects or rodents.
- Never deliver food in a vehicle used to haul garbage.
- Do not bring pets when delivering food.
- Keep items that could contaminate food separate from the delivery. This could include oil, antifreeze, or wiper fluid.
- Lock and seal delivery vehicles when they are not being loaded or unloaded.

It is critical that food remain at safe temperatures during transport. If not, bacteria on the food can grow and cause an illness. Here's what you should do to prevent this.



- Check the temperature of refrigerated trucks to make sure they can keep food at 41°F (5°C) or lower.
 - Check the readout on the truck's temperature recording and monitoring device. Document the reading.
 - [•] It is recommended that refrigeration units remain on and vehicle doors remain closed during periods of inactivity.
 - [°] Refrigerated trucks must be pre-cooled prior to loading.
- Make sure coolers, ice packs, and thermal blankets are used in unrefrigerated vehicles to keep cold food cold.

Preparing Food for Transport

When getting food ready for delivery, follow these guidelines to keep it safe.



DO NOT leave food outside and unsupervised.



DO NOT put food by garbage containers.



Use boxes that are sturdy, clean, and dry.



DO NOT leave refrigerated or frozen food at room temperature.



Keep shelf-stable food in a clean, dry storage area.



Keep food in a pest-free area.



Keep meat, poultry, and seafood separate from each other and away from other food.



Keep chemicals separate from food.

Loading and Transporting Food Safely

Food must be loaded into the delivery vehicle correctly to prevent contamination. Once loaded, food must stay at the correct temperature to keep it safe. Follow these guidelines when loading and transporting food.



- Keep refrigerated food at 41°F (5°C) or lower during transport.
- If possible, keep frozen food at temperatures that will keep it frozen.



- Keep food cold in unrefrigerated vehicles.
 - Always cover refrigerated and frozen food with thermal blankets.
 - Or place it in coolers with ice packs.
- DO NOT store allergens over other products.



- DO NOT store raw food over readyto-eat food.
 - Raw food includes raw meat, seafood, poultry, and shell eggs.



- DO NOT store chemicals with food products.



and frozen food so air can circulate around it. ^o Use pallet slip

sheets.

Load refrigerated



 Keep drive times in unrefrigerated vehicles to 30 minutes or less.



- Check the temperature of refrigerated food when it has reached its destination.
- Document the temperatures.

Apply Your Knowledge

Was it Loaded Safely?

ullet Write an X next to each situation where a mistake was made.



Notes

- 24

-

_

10



- How and when to clean and sanitize
- How to manually clean and sanitize tools and utensils
- Handling garbage
- Spotting pests



Pest Troubles

An agency was closed by the local health department after officials found significant evidence of rodents in the operation. Droppings were spotted in food-storage areas and nests were found near walls. Many food products were also damaged by the rodents. Aside from the damage, officials feared that pathogens carried by the pests would make clients sick. The health department worked closely with the agency to implement pest prevention measures including correct cleaning and sanitizing procedures.

HOW AND WHEN TO CLEAN AND SANITIZE

Cleaning removes food and other dirt from a surface. Sanitizing reduces pathogens on a surface to safe levels. The most important reason to clean and sanitize is to prevent the spread of pathogens to food. It also helps control pests such as insects and rodents.

Surfaces to Clean and Sanitize

Not all surfaces are handled the same way. Some only need to be cleaned. Others must be cleaned and sanitized.



- All surfaces must be cleaned and rinsed. Examples include:
- Walls and floors
- Storage shelves
- Garbage containers



Any surface that touches food must be cleaned and sanitized. Examples include:

- Plastic food bins
- Ladles
- Scales
- Prep tables in clean rooms
- Sorting tables
- Scoops



If you notice worn or cracked equipment, set it aside and report it to your manager.

This equipment is not easy to clean or sanitize. It also may hold pathogens.

How to Clean and Sanitize Surfaces

The procedure below shows you how to clean and sanitize. Be sure to avoid contamination when cleaning. For example, do not allow spray cleaner to contaminate food. Always use cleaners and sanitizers according to manufacturers' directions and your organization's policies.

Scrape or remove food from the surface.





Wash the surface.



Rinse the surface.



Sanitize the surface.

5

Allow the surface to air-dry.

Cleaning and Sanitizing Stationary Equipment

Cleaning stationary equipment is similar to cleaning other surfaces. Talk to your director or supervisor about how to clean specific equipment you might use. Follow these steps when cleaning and sanitizing stationary equipment.



- Unplug the equipment.
- Take off the parts that can be removed. Wash, rinse, and sanitize them by hand.



- Scrape or remove food from the equipment surfaces.
- Wash, rinse, and sanitize the equipment surfaces.
- Let the surfaces air-dry. Then put the equipment back together.

When To Clean and Sanitize

All food-contact surfaces need to be cleaned and sanitized at these times.



Before and after use



 Before repacking food



• When changing to a new product or between allergens



- After four hours if the items have been in constant use
- Any time the item has become contaminated

How to Make Sure Sanitizers Are Effective

You must make sure the sanitizer you use is effective. Ask your director or supervisor how to do these steps.



• Make sure the water is the correct temperature for the sanitizer you are using.



- Make sure you have used the correct amount of sanitizer.
- Use a test kit to check the sanitizer's strength.



- Make sure you leave the items being sanitized in contact with the sanitizer for the correct amount of time. This will reduce pathogens to safe levels.
- When wiping or spraying sanitizer on surfaces such as prep tables or sorting tables, allow the sanitizer to air-dry on the surface.

Apply Your Knowledge Cleaning and Sanitizing

 $oldsymbol{1}$ Circle the item that needs to be both cleaned and sanitized. $oldsymbol{1}$





Cleaning and Sanitizing continued

Apply Your Knowledge

Put the steps for cleaning and sanitizing in the correct order by placing the number of each step in the space provided.			
A	Sanitize the surface.		
B	Wash the surface. Allow the surface to air-dry. Rinse the surface.		
C			
D			
E	E Scrape or remove food from the surface.		
Write the ite	an $ imes$ next to each situation that requires the food handler to clean and sanitize em being used.		
A	Jorge has used the same scoop to portion cereal for an hour.		
B	Bob drops the scoop on the floor while using it.		
C	Shelly has finished repacking produce on a prep table and now will use it to repack cereal.		
D	Maria is about to start scooping bulk rice into smaller bags.		
Wher unplu	n cleaning and sanitizing stationary equipment, what is the next step after gging it?.		
A	Taking off any removable parts and washing, rinsing, and sanitizing them		
B	Scraping or removing food from the equipment surfaces		
C	Washing, rinsing, and sanitizing the stationary equipment surfaces		
D	Letting all surfaces air-dry		
9 Write	a an \times next to the actions that prevent a sanitizer from working well.		
A	Making the sanitizer temperature too high		
B	Putting extra sanitizer in the solution		
C	Letting items make contact with the sanitizer solution		
D	Testing the sanitizer strength with a test kit		

How to MANUALLY CLEAN AND SANITIZE TOOLS AND UTENSILS

Tools and utensils are usually cleaned and sanitized in a three-compartment sink. The sink must be set up the correct way before it can be used. Once set up, there are specific steps you must follow to make sure items are cleaned and sanitized correctly.

Setting Up a Three-Compartment Sink

Start by cleaning and sanitizing each sink and all work surfaces. Next set up each sink following these steps.



Sink 1

- Fill with water at least 110°F (43°C).
- Add detergent. Ask your director or supervisor how to do this.



Sink 2

• Fill with water. Leave the sink empty if you spray-rinse items.



Sink 3

- Fill with water.
- Add the correct amount of sanitizer. Ask your director or supervisor how to do this.
- Check the strength of the sanitizer with a test kit.

Using a Three-Compartment Sink

Follow these steps to clean and sanitize items in a three-compartment sink.



Rinse, scrape, or soak the items before washing them.



Wash the items in the first sink.

- Use a brush, cloth, or nylon scrub pad to loosen dirt.
- Change the water and detergent when the suds are gone or the water is dirty.



Rinse the items in the second sink.

- Dip them in the water or spray-rinse them.
- Remove any food or detergent.
- Change the water when it becomes dirty or full of suds.



Sanitize the items in the third sink.

- Soak them in the sanitizer solution for the correct length of time.
- **NEVER** rinse items after sanitizing them. This could contaminate the surfaces.



Air-dry the items.

- Place them upside down so they will drain.
- Do NOT wipe them dry.



Washing Tools and Utensils

- 1 Write an × next to the first thing you should do when setting up a three-compartment sink.
- _____ A Fill the sinks with water.
- _____ B Mix the sanitizing solution.
- _____C Mix the detergent solution.
- _____D Clean and sanitize the sinks.

strate and containants are and containents a

2 Put the steps for cleaning and sanitizing items in a three-compartment sink in order.

- _____ A Air-dry the items.
- _____ B Rinse the items.
- _____ C Sanitize the items.
- _____ D Rinse, scrape, or soak the items.
 - E Wash the items.

HANDLING GARBAGE

Garbage can contaminate food and equipment if it's not handled safely. It can also create odors and attract pests. Here's how you should handle it.



- Remove garbage as quickly as possible.
- Be careful not to contaminate food or surfaces when removing garbage.



- DO NOT clean garbage containers in clean rooms or food-storage areas.
- Clean the inside and outside of garbage containers often.



- Close the lids on outdoor containers.
- Keep indoor food waste containers covered.



That's Just Garbage

1 Draw an \times through each situation that is unsafe.





SPOTTING PESTS

Pests, such as insects and rodents, carry pathogens that can make people sick. Insect parts can even be a source of physical contamination. That's why it is important to look for signs that pests are in the food bank or agency. Tell your director or supervisor if you spot these signs.



Droppings



Nests

• Damage to products, packaging, and the facility



Notes	
-------	--



÷.

 \mathbf{x}_{i}

.

2

What Do You Think?

- A. Leaving raw chicken on a pallet in an unrefrigerated loading area is time-temperature abuse. This can allow pathogens on the chicken to grow.
- B. It is poor personal hygiene to sneeze on food. This can transfer pathogens from the body to food.
- C. Allowing raw meat to drip juices onto produce is cross-contamination. This can allow the transfer of pathogens on the meat to the produce.
- D. The food-storage container was not cleaned and sanitized correctly. Pathogens could be transferred to food stored in the container.

The Most Common Food Allergens

2; 3; 5; 6; 10; 14; 16; 17

Check Your Handwashing Knowledge

A. Ryan applied hand sanitizer before washing his hands. This should only be done after handwashing.

D. Ryan only scrubbed his hands with soap for five seconds. He should have scrubbed his hands with soap for at least 10 to 15 seconds. F. Ryan dried his hands with his apron. Hands should only be dried with single-use lowels or a hand dryer.

H. Ryan opened the restroom door with his clean hands, possibly contaminating them again. He should have opened the restroom door using a paper towel.

- A; B; D
- B C

Keeping It Safe

B. Alicia should not have rinsed her gloves. Food handlers must never rinse, wash, or reuse gloves.

C. Alicia should not have repacked rice after repacking cereal without changing her gloves. Gloves must always be changed in-between tasks.

B. It is unsafe to reuse the same pair of single-use gloves. Gloves must never be reused. They must also be changed in-between tasks

C. Prepping food while wearing false fingernails is unsafe. Food handlers with nail

polish or false fingernails must wear gloves when handling exposed food D. It is unsafe to work with an infected cut that is not bandaged and covered correctly. Infected wounds on the hand or wrist must first be covered with a bandage or finger cot that will prevent fluids from leaking out. Then a single-use glove must be worn over top.

It's What You Wear

- Wearing a dirty shirt is unsafe because pathogens on it could contaminate exposed food.
 - 2. Wearing nail polish without wearing single-use gloves is unsafe. Nail polish can hide dirt under nails and may flake off into exposed food.
 - 4. Wearing watches can be unsafe when handling exposed food or when working around it. Watches, like other jewelry on the hands and arms, can contain bacteria and other pathogens. These pathogens could get onto the exposed food.
 - 5. Touching an apron when removing it can contaminate it. Disposable aprons must be thrown away when leaving the clean room. A new apron should be put on before returning to the clean room.
 - 7. Cleaning and sanitizing utensils and equipment used to handle exposed food without wearing a hair covering can contaminate it. Always wear a clean hat or other hair covering when working in areas used to clean utensils and equipment.
 - 8. Wearing the same pants day after day can be unsafe because they can pick up pathogens. Wear clean clothing whenever working with or around food.

Reporting Illnesses

1; 2; 3; 5

Ice-Point Calibration Method

1 D

2 D

Is the Delivery Safe?

A. A broken door lock on a delivery vehicle is unsafe. This can allow tampering with the delivery. Door locks must be in place and functioning on delivery vehicles.

B. Any sign of pests in a delivery vehicle is

unsafe. A food delivery should be rejected if delivery containers or the delivery vehicle contains pests.

D. Homemade products, such as the home-canned jelly, may be unsafe. You should not receive these products.

Is It the Correct Temperature?

A. The frozen meat was not received at the correct temperature. It will start to thaw at 38°F (3°C). Frozen meat must be received frozen solid.

B. The bags of cut lettuce where not received at the correct temperature. Cut produce must be received at 41°F (5°C) or lower.

C. Cut melons received at 45°F (7°C) may be unsafe. Cut produce must be received at 41°F (5°C) or lower.

D. Milk received at 50°F (10°C) may be unsafe. Refrigerated food, such as milk, should be received at 41°F (5°C) or lower unless otherwise specified by the manufacturer.

Is It Stored Correctly?

 A. This food has not been stored correctly. Food must be stored away from walls.
 B. The sour cream is not being stored at the correct temperature. Refrigerated food must be stored at 41°F (5°C) or lower.

D. This food is being stored on the floor, which is not correct. Food must be stored at least six inches (15 cm) off the floor.

Has it Been Rotated Correctly?

в в

Keep It or Toss It?

- 1. This jar has a severe dent in the lid. It should be rejected.
 - 2. This can has a deep dent in the can body. It should be rejected.
 - 3. This jar has a missing label. It should be rejected.
 - 5. This can has a bulging end. It should be rejected.
 - 7. This jar is cracked. It should be rejected.
 - 8. This canned food is homemade. It should be rejected.

Should You Keep It?

- Some of these oranges are moldy. They should be rejected.
 - 3. The skin on these tomatoes is not intact. They should be rejected.
 - 4. This pasta is exposed. It should be rejected.

Is It Acceptable?

- I his medication has a broken tamper-resistant seal. It should be rejected.
 - 2. This medication is expired. It should be rejected.
 - 3. This medication has a label that is not readable. It should be rejected.

What Did They Do Wrong?

- 1. These scoops are not being stored correctly. They may become contaminated.
 - 2. These people are repacking two different food products at the same time. This can be unsafe.
 - 3. This person is repacking exposed food without wearing a hat or other hair covering. This is unsafe.

Was it Loaded Safely?

- 1. Eggs were loaded on top of other food in this delivery vehicle. This is unsafe because eggs, an allergen, are being stored over other food, which can result in contamination.
 - 2. Chemicals were loaded on the same pallet with food. Chemicals should never be stored with food because they can contaminate it.

Cleaning and Sanitizing

- B. A scale used in a clean room to weigh exposed food must be cleaned and sanitized.
- A. 4; B. 2; C. 5; D. 3; E. 1
- B. The scoop must be cleaned and sanitized because it may have become contaminated after falling on the floor.

C. The prep table must be cleaned and sanitized. Items that will come in contact with exposed food must be cleaned and sanitized when changing to a new product or between allergens.

D. The scoop that Maria will be using must be cleaned and sanitized. Items that will come in contact with exposed food must be cleaned and sanitized before and after use.

I A

A, B

Washing Tools and Utensils

🖲 D

A. 5; B. 3; C. 4; D. 1; E. 2

That's Just Garbage

 Carbage should not be stored against a wall in the facility. It should be removed from the facility as quickly as possible.

TEST

Circle the best answer to each question below.



The three types of hazards that make food unsafe are

- A biological, chemical, and physical.
- B bacterial, nutritional, and universal.
- C fungus, viruses, and parasites.
- D allergens, sanitizers, and contaminants.

Transferring pathogens from the body to food can be prevented by practicing correct

- A cleaning and sanitizing.
- B temperature control.
- C personal hygiene.
- D receiving and storage.
- A food allergy occurs when the body has a reaction to
 - A cleaning chemicals.
 - B airborne pathogens.
 - C undercooked meat.
 - D certain food items.
- Which is a common food allergen?
 - A Sugar
 - B Lettuce
 - C Tomatoes
 - D Wheat

What occurs when food containing an allergen comes in contact with other food?

- A Cross-contamination
- B Cross-contact
- C Biological contamination
- D Biological contact

- One way to keep people with food allergies safe is to
 - A freeze food before distributing it.
 - B prevent cross-contact between food items.
 - C store food with allergens on top of other food.
 - D wipe off open food that has touched an allergen.
- 2

Hands and arms should be scrubbed with soap for how many seconds during handwashing?

- A 1-3
- B 4-6
- C 7-9
- D 10-15
- 8 When should hands be washed?
 - A After putting on gloves
 - B After touching the hair or body
 - C Before taking out garbage
 - D Before sneezing, coughing, or using a tissue

What activity can be performed at a handwashing sink?

- A Dumping mop water
- B Rinsing food scoops
- C Cleaning canned food
- D Washing hands only

- How should hair be worn when repackaging food?
 - A Clipped back with hair pins
 - B Tied back with a rubber band
 - C Pulled off the face with a headband
 - D Covered with a hat or other hair covering
- Eating and drinking when working in food-storage areas is a food safety hazard because it could
 - A cause choking.
 - B offend clients.
 - C contaminate hands and food.
 - D lead to time-temperature abuse.

¹⁵ What symptom must be reported to the director or supervisor?

- A Headache
- **B** Dizziness
- C Diarrhea
- D Sweating

The temperature danger zone is the temperature range between

- A 0°F to 41°F (-18°C to 5°C)
- B 32°F to 41°F (0°C to 5°C)
- C 41°F to 135°F (5°C to 57°C)
- D 135°F to 212°F (57°C to 100°C)

¹⁴ A delivery should be rejected when

- A the truck contains a door lock.
- B the product contains pests.
- C refrigerated food is 41°F or lower.
- D frozen food is frozen solid.



A bimetallic stemmed thermometer is being calibrated. After the stem has been submerged in ice water and the indicator has stopped moving, the thermometer must be adjusted to what temperature?

- A 0°F (-18°C)
- B 32°F (0°C)
- C 41°F (5°C)
- D 45°F (7°C)

15 Refrigerated food must be stored at temperatures no higher than

- A 41°F (5°C).
- B 45°F (7°C).
- C 47°F (8°C).
- D 49°F (9°C).
- How far off the floor must food be stored?
 - A 1 in (3 cm)
 - B 2 in (5 cm)
 - C 4 in (10 cm)
 - D 6 in (15 cm)
- ¹⁸ Which storage date is the last date recommended for the product while at peak quality?
 - A Packing or manufacturing date
 - B Sell-by date
 - C Best-by or Best If Used By date
 - D Use-by or expiration date
- ¹⁹ Cans should be discarded if they have
 - A torn labels that are still readable.
 - B shallow dents in the can body.
 - C swollen or bulging ends.
 - D rust that can be wiped off.



20 Produce should be discarded when it has

- A mold.
- B light bruising.
- C code dates.
- D intact skin.
- TCS food should be repackaged in a clean room that maintains a temperature no higher than
 - A 55°F (13°C)
 - B 65°F (18°C)
 - C 75°F (24°C)
 - D 85°F (29°C)
- 23 What information must be included on repackaged food labels?
 - A Nutritional value of each ingredient
 - B Directions for preparation
 - C List of each major food allergen
 - D Calorie count of each serving

Drive times in unrefrigerated delivery vehicles should be kept to less than

- A 120 minutes.
- 90 minutes. B
- C 60 minutes.
- D 30 minutes.
- Which practice when transporting food will help keep it safe?
 - A Keeping refrigerated food at 45°F or lower
 - B Storing raw food over ready-to-eat food
 - C Packing the delivery vehicle tightly so air cannot circulate around products
- D Covering refrigerated food with thermal blankets

- ²⁵ What is the difference between cleaning and sanitizing?
 - A Cleaning removes dirt while sanitizing reduces pathogens to safe levels
 - B Cleaning removes dirt and pathogens while sanitizing sterilizes a surface
 - C Cleaning reduces pathogens to safe levels while sanitizing removes visible dirt and food
 - D Cleaning reduces pathogens to safe levels while sanitizing sterilizes a surface
 - What is the correct order for cleaning and sanitizing?
 - A Rinse, wash, sanitize, scrape, air-dry
 - B Rinse, sanitize, wash, scrape, air-dry
 - C Scrape, rinse, wash, sanitize, air-dry
 - D Scrape, wash, rinse, sanitize, air-dry
 - What surface must be cleaned and sanitized?
 - A Refrigerator door
 - B Prep table
 - C Storage shelf
 - D Garbage container
 - What is the second sink of a three-compartment sink used for?
 - A Scraping
 - B Washing
 - C Rinsing
 - D Sanitizing
 - Where should garbage containers be cleaned?
 - A In clean rooms
 - B In three-compartment sinks
 - C In salvage areas
 - D Away from food-storage areas



30 Insects and rodents are a food-safety risk because they

- A eat food that is meant for clients.
- B carry pathogens and can make people sick.
- C scare employees.
- D make a mess with their nests and droppings.



) (2) 800.765.2122 servicecenter@restaurant.org

Distributed by



175 W Jackson 8171, Ste 1500 Chicago IL 60604-2814 1.800.765 2122 Restaurant.org | Serv Safe.com

 2014 National Restaurant Association Educational Foundation. All rights reserved. ServSale - and the ServSale rogo are trademarks of the NRAEF. National Restaurant Association - and the arc design are trademarks of the National Restaurant Association. Feeding America: All rights reserved. Feeding America (s a SO1 (c)/(3) non-profit recognized by the IRS.



